

Dicke model: entanglement as a finite size effect

Oleksandr Tsyplatyev and Daniel Loss

Department of Physics, University of Basel, Klingelbergstrasse 82, CH-4056 Basel, Switzerland

(Dated: July 15, 2009)

We analyze the Dicke model at zero temperature by matrix diagonalization to determine the entanglement in the ground state. In the infinite system limit the mean field approximation predicts a quantum phase transition from a non-interacting state to a Bose-Einstein condensate at a threshold coupling. We show that in a finite system the spin part of the ground state is a bipartite entangled state, which can be tested by probing two parts of the spin system separately, but only in a narrow regime around the threshold coupling. Around the resonance, the size of this regime is inversely proportional to the number of spins and shrinks down to zero for infinite systems. This spin entanglement is a non-perturbative effect and is also missed by the mean-field approximation.

Coherent interaction between electromagnetic photon fields and matter attracted interest a long time ago [1] with renewed attention gained in the last decade due to significant developments in the experimental techniques in various areas of physics. Achievement of Bose Einstein condensation of cold atomic gases in electromagnetic traps enabled the coherent coupling of hyperfine states of 10^5 atoms to a single photon mode of an optical resonator [2]. Advances in the semiconductor technology allowed to obtain optical microcavities where electron-hole excitations inside the semiconductor quantum well are strongly coupled to an eigenmode of the optical resonator [3]. Strong coupling of a single mode of a transmission line resonator to a Cooper pair box [4] and a quantized mode of an optical crystal cavity to several semiconductor quantum dots [5] have been demonstrated as a possible way to a quantum computing device [6].

Theoretical understanding of all these systems is based on a model proposed by Dicke [7] which describes N spins $1/2$ (identical two-level systems) with splitting energy 2ϵ coupled to a single mode of electromagnetic field ω . It was shown that this model is exactly diagonalisable [8]. At zero temperature it undergoes a quantum phase transition from a non interacting state with unpopulated bosonic mode to a condensed state with a highly populated bosonic mode [9] if coupling between the boson and a single spin g is greater than a threshold value. In the thermodynamic limit a phase transition occurs in the region of strong coupling if temperature is less than a critical temperature which can be described by a Bogolyubov Hamiltonian similarly to the pairing model of superconductivity [10]. Recently, a variational wave function approach to the generalised Dicke model was used [11] to describe Bose Einstein condensation of exciton-polaritons in a semiconductor optical cavity.

In this paper we analyse the Dicke model at zero temperature for a finite N using matrix diagonalisation methods. We find that for the particular coupling strength,

$$\omega\epsilon < g^2 \lesssim \omega\epsilon \left(1 + \frac{1}{N}\right), \quad (1)$$

the ground state of the spin subsystem is a bipartite entangled state and it is not entangled outside of this re-

gion. The lower bound of the inequality is onset of the quantum transition described by the mean field theory [12]. The upper bound is the condition to have only singly populated bosonic mode in the ground state. The approximated value is the result of $1/N$ expansion around the resonance.

In the thermodynamic limit the ground state projected onto the subspace of N spins is not entangled as it is a product state. For a weak coupling below the quantum transition the ground state is a product of all unexcited single spin states. For a strong coupling above the transition threshold the ground state is also a product state of all single spin states [11] as a result of the mean-field approximation. In a finite system the mean-field approximation is not applicable in a small region above the transition threshold where the expectation value of the boson is of the order of 1 and its fluctuations are also of the order of 1. The ground state in this region is a superposition of the unexcited state of all spins and a spin state with only one single spin-flip excitation, the N spin W-state, $|W\rangle = (|\uparrow\downarrow\downarrow\dots\rangle + |\downarrow\uparrow\downarrow\dots\rangle + |\downarrow\downarrow\uparrow\dots\rangle + \dots)/\sqrt{N}$. The W-state can also be interpreted as 'magnon state' at vanishing wave vector [13].

Furthermore, the W-state can be considered as a bipartite entangled state in the following sense. Dividing all spins into two groups [14] the W-state is a Bell state in the subspace restricted by only one spin-flip excitation above the unexcited states of each group of the spins, see Eq. (13). In the course of a bipartite measurement if the first group is found in the excited state then the second group is projected onto the unexcited state and if the first group is found in the unexcited state then the second group is projected onto the excited state.

We diagonalize the Dicke model for N spins $1/2$ coupled to a single bosonic mode

$$H = \omega b^\dagger b + \epsilon \sum_j S_j^z + \frac{g}{\sqrt{N}} \sum_j (S_j^+ b + S_j^- b^\dagger), \quad (2)$$

where the sum runs over N spin- $1/2$ operators $\mathbf{S}_j$ that obey the commutation relations $[S_i^\alpha, S_j^\beta] = \epsilon_{\alpha\beta\gamma} \delta_{ij} S_i^\gamma$, and b ($b^\dagger$) is standard bosonic annihilation (creation) operator.

The Dicke model possesses the following conserved quantities. One is the number of excitations of the coupled spin-boson system,

$$L = n + J_z, \quad (3)$$

expressed in terms of the z-component of total spin operator $J_\alpha = \sum_j S_j^\alpha$, $\alpha = x, y, z$, and the occupation number operator $n = b^\dagger b$ of the boson mode. Note that J_α and n are not conserved separately. The eigenvalues of L are the so-called cooperation numbers c , given by the sum of expectation values of n and J_z . A second conserved quantity is the total spin, $J^2 = J_z^2 + (J_+ J_- + J_- J_+)/2$ with eigenvalues $j(j+1)$.

We represent the Dicke Hamiltonian (2) in a basis where J^2 and L are block-diagonal. Within a block of given c and j the remaining degrees of freedom can be labeled by the eigenvalues m of J_z . In the representation $|c, j, m\rangle$ each block has a tridiagonal form. The diagonal matrix elements represent the energies of states containing $c - m$ bosons and m excited spins,

$$\langle m|H|m\rangle = \omega(c - m) + \epsilon m. \quad (4)$$

The first above- and below-diagonal matrix elements are transition amplitudes connecting all pairs of states which differ by just one flipped spin,

$$\langle m|H|m+1\rangle = \frac{g}{\sqrt{N}} \sqrt{(c - m)(j(j+1) - m(m+1))}. \quad (5)$$

The size of each block is limited by the fact that $-N/2 \leq m \leq N/2$ for $j = N/2$. However, the upper bound on m is further constrained by the cooperation number c which has the lower bound $-N/2$ for a block with no bosons present in the state.

The ground state of Eq. (2) is the lowest energy state of all blocks Eqs. (4,5). For different values of the parameters ω , ϵ , and g the ground state can have different c 's and j 's (see below).

We consider now the case with the bosonic (ω) and spin excitation energies (ϵ) close to each other. For the uncoupled case, $g = 0$, the ground state contains no bosons and no excited spins (i.e. all spins are, say, down). From Eqs. (3) and (4) we see that the cooperation number of this state is then $c = -N/2$ with corresponding ground state energy $E_0 = -N\epsilon/2$. For finite but still small coupling, $g \ll 1$, we can use a perturbative approach to remove the boson mode via a Schrieffer-Wolff transformation to obtain an effective spin-Hamiltonian [16],

$$H = \omega b^\dagger b + \left(\epsilon + \frac{2g^2 b^\dagger b}{N(\epsilon - \omega)} \right) \sum_j S_j^z + \frac{g^2}{N(\epsilon - \omega)} \sum_{i,j} S_i^+ S_j^-, \quad (6)$$

where the spin-boson mixing is eliminated up to the second order in g [15], introducing an effective XY coupling




Figure 1: Cooperation number c of the ground state obtained by matrix diagonalization of the Dicke model for $N = 3$. The first solid line from the left indicates the quantum transition to the strongly correlated non-perturbative regime, and the second solid line is determined by Eq. (9). The dashed line corresponds to the resonance $\epsilon = \omega$.

between the spins within a subband of given boson occupation number n (with n being conserved under the effective Hamiltonian Eq. (6)). Such a perturbative approach gives a qualitatively correct description of the energies and wave functions until g crosses a threshold value g_c where a transition to a strongly correlated non-perturbative regime takes place (see below). The cooperation number c of the ground state for the Dicke Hamiltonian Eq. (2) is plotted in Fig.1 for the solution for $N = 3$. There are several regimes: for small g the ground state of the system is defined by a regime (black) with $c = -N/2$ and $\langle n \rangle = 0$. Then, with increasing coupling a quantum phase transition at g_c takes place to a new regime with $c > -N/2$ where $\langle n \rangle > 0$, and where subbands with sharp bosonic occupation numbers no longer exist.

Increasing g further above the transition threshold a sequence of states with $c = 1 - N/2$, $c = 2 - N/2$, and so on becomes subsequently the ground state due to the interaction energy. The ground state with c greater but close to $-N/2$ can not already be approximated with help from the perturbation theory as coupling is too strong and can not yet be approximated using the mean field approach as the fluctuations of the order parameter are too large compared to its mean. Here we diagonalize the Hamiltonian Eq. (2) using the matrices Eqs. (4,5). Sizes of the matrices in this regime are limited by the cooperation number c . In the presentation $|c, j, m\rangle$ the matrices are not larger than 2×2 and 3×3 for $c = 1 - N/2$ and $c = 2 - N/2$ respectively. The lowest eigenenergies of these matrices are

$$E_1 = -\frac{(N-1)\epsilon}{2} + \frac{\omega}{2} - \sqrt{g^2 + \left(\frac{\omega - \epsilon}{2}\right)^2} \quad (7)$$

for $c = 1 - N/2$ and

$$E_2 = -\frac{(N-2)\epsilon}{2} + \omega - 2g\sqrt{1 - \frac{1}{2N}} + \frac{\omega - \epsilon}{2(2N-1)}. \quad (8)$$

for $c = 2 - N/2$. The last expression is a result of expansion at the resonance in powers of $\omega - \epsilon$. Both E_1 and E_2 belong to subblocks with $j = N/2$ in accordance with a theorem from [10]. Comparing E_0 and E_1 we find $g_c = \sqrt{\omega\epsilon}$. The same condition to have a non zero population of the bosonic mode in the ground state was established in [9].

Comparing E_1 and E_2 near the resonance $\omega = \epsilon$ we find that a state with $c = 2 - N/2$ becomes the ground state when g exceeds some value g_2 given by

$$g_2 = \frac{\epsilon + \omega + (\omega - \epsilon) / (2N - 1)}{2 \left(2\sqrt{1 - 1/2N} - 1 \right)}. \quad (9)$$

Thus, $g < g_2$ together with $g > g_c$ define the regime of the model parameters where matrix diagonalization is the only way to study the Dicke model. The upper bound of Eq. (1) coincides with Eq. (9) for $\epsilon \approx \omega$ and $N \gg 1$. Increasing g further, we can determine the boundaries between the ground states with different values of c (i.e. $c = 3 - N/2, 4 - N/2, \dots$) by numerical diagonalization, as shown in Fig.1 for $N = 3$. A result similar to Eqs. (7, 8, 9) was obtained in [17] but the discrete jumps in the cooperation number of the ground state were interpreted as an infinite sequence of instabilities.

In the strong coupling regime, $g \gg g_c$, the mean field approach provides a good approximation to the exact ground state. Indeed, introducing the expectation value of the bosonic operator $\mathcal{B} = \langle b \rangle$ and neglecting quantum fluctuations around it, the Dicke Hamiltonian Eq. (2) becomes

$$H = \omega |\mathcal{B}|^2 + \sum_j \left(\epsilon S_j^z + \frac{g}{\sqrt{N}} (S_j^+ \mathcal{B} + S_j^- \mathcal{B}^*) \right). \quad (10)$$

Eigenstates of this Hamiltonian are product states of the N spins, and thus are manifestly not entangled. Diagonalization of the 2×2 - matrices for each spin and subsequent minimization of the sum of the lowest eigenenergies over $|\mathcal{B}|$ gives the following self-consistency (mean-field) equation

$$\omega = \frac{g^2}{\sqrt{4|\mathcal{B}|^2 g^2 / N + \epsilon^2}}, \quad (11)$$

which describes a quantum phase transition at a threshold value of the coupling strength g_c that we have already found from the matrix diagonalization.




Figure 2: Expectation value of n calculated by matrix diagonalization of Eqs. (4,5) - solid line and the mean field result - dashed line. Plots are for $N = 3$ spins and at resonance $\epsilon = \omega$. The grey area is the regime defined by Eq.(1) where the cooperation number of the ground state is $c = 1 - N/2$.

The mean field approximation is satisfactory even for a system of only a few spins. To see this, we compare for $N = 3$ spins in Fig.2 the expectation value of n taken with respect to the exact ground state of Eqs. (4,5) with $|\mathcal{B}|^2$ given in Eq. (11) at the resonance $\omega = \epsilon$. There we see that the largest deviation is in the region of intermediate coupling strength where the cooperation number of the exact ground state is $c = 1 - N/2$, grey area in Fig.2. For weak coupling $g < g_c$ the ground state coincides with the non-interacting one and $\langle n \rangle = |\mathcal{B}|^2 = 0$. The quantum fluctuations of n compared to its mean value are already small for the coupling corresponding to the exact ground state with $c = 4 - N/2$, $(\langle n^2 \rangle - \langle n \rangle^2) / \langle n \rangle^2 \simeq 0.2$. Thus, in the strong coupling regime the approximation of neglecting these fluctuations in Eq. (10) is already good for $g \gtrsim g_2$ irrespective of N .

The ground state can be characterized in terms of entanglement between different parts of the spin-subsystem. In the weak coupling regime, the ground state, being a direct product of (unexcited) spin states, is not entangled. In the regime of strong coupling, where the approximate Hamiltonian Eq. (10) is valid, the ground state is also a product of the individual spin states which, thus, also has no entanglement of any pair of spins. The ground state in the intermediate region Eq. (1) has to be found by matrix diagonalization and will be analyzed below.

Diagonalization of the matrix Eq. (4,5) for $c = 1 - N/2$ and $g = \sqrt{\omega\epsilon}$ gives a ground state in the regime defined by Eq. (1). We change the mixed spin-boson representation $|c, l, m\rangle$ to a separate representation of spins and the boson $|\mathbf{n}, l, m\rangle$, where $\mathbf{n}$ is the bosonic occupation number. Then, tracing out the bosonic degree of freedom $\mathbf{n}$

we obtain the reduced density matrix of the spins only, which in the representation of J_z eigenstates $|m\rangle$ is given by

$$\hat{\rho} = \frac{\epsilon}{\epsilon + \omega} \left| -\frac{N}{2} \right\rangle \left\langle -\frac{N}{2} \right| + \frac{\omega}{\epsilon + \omega} \left| -\frac{N}{2} + 1 \right\rangle \left\langle -\frac{N}{2} + 1 \right|. \quad (12)$$

There is a finite probability to find either all spins in the completely polarized state or in the W-state $| -N/2 + 1 \rangle$.

The W-state is a Bell state for $N = 2$. For $N > 2$ it is a bipartite entangled state. The set of spins can be divided into two equal groups consisting of $N/2$ spins each (assuming N even). In the basis $|m_1\rangle |m_2\rangle$, m_1 and m_2 are the eigenvalues of the operators $J_z^{1(2)}$ belonging to the first (second) group, the state $| -N/2 + 1 \rangle$ is

$$\left(\left| -\frac{N}{4} \right\rangle \left| -\frac{N}{4} + 1 \right\rangle + \left| -\frac{N}{4} + 1 \right\rangle \left| -\frac{N}{4} \right\rangle \right) / \sqrt{2}, \quad (13)$$

i.e. a measurement of one group projects another group onto the definite state. Therefore in the intermediate region Eq.(1) the ground state of the Hamiltonian Eq. (2) is a bipartite entangled state. Note that N spin W-state does not belong to the class of N spin entangled states prepared by squeezing [18]. The squeezing parameter is

not defined for the $N = 2$ W-state and is greater than one for $N > 2$ W-states.

Let the spins have different splitting energies ϵ_j in Eq.(2) instead of the case $\epsilon_j \equiv \epsilon$ which we have considered in the paper. A small variation $\langle \epsilon_j \rangle - \epsilon_j \ll \epsilon_j$, $\langle \epsilon_j \rangle$ being the average over splitting energies of all spins, can be treated using perturbation theory and will not affect our results, Eqs.(1,13) much. For instance, the $N = 2$ W-state in Eq.(13) becomes $((1 - (\epsilon_1 - \epsilon_2)/\sqrt{2}g)|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle)/\sqrt{2}$ at the resonance $\epsilon = \omega$. The amplitudes of different components in the singlet will alter slightly while the W-state will not change qualitatively. For a more detailed analysis of the inhomogeneous Dicke model see [19].

In conclusion, we analyzed the Dicke model for a finite-size system by matrix diagonalization. We found that the ground state is a bipartite entangled state only in a narrow regime of parameters next to the quantum phase transition at zero temperature. The ground state in this regime cannot be obtained from mean-field theory which approximates the ground state as a product state. Also perturbation theory is not valid in this regime. For an infinite system width of the corresponding parametric region vanishes as $1/N$ if system is close the resonance.

We acknowledge financial support from the Swiss NF, NCCR Nanoscience Basel, and JST ICORP.

-
- [1] C. W. Gardiner and P. Zoller, *Quantum Noise*, Springer, 2004.
 - [2] F. Brennecke, T. Donner, S. Ritter, T. Bourdel, M. Köhl, and T. Esslinger, *Nature* **450**, 268 (2007).
 - [3] J. Kasprzak, M. Richard, S. Kundermann, A. Baas, P. Jeambrun, J. M. J. Keeling, F. M. Marchetti, M. H. Szymańska, R. André, J. L. Staehli, V. Savona, P. B. Littlewood, B. Deveaud, and Le Si Dang, *Nature* **443**, 409 (2006).
 - [4] A. Wallraff, D. I. Schuster, A. Blais, L. Frunzio, R.-S. Huang, J. Majer, S. Kumar, S. M. Girvin and R. J. Schoelkopf, *Nature* **431**, 162 (2004).
 - [5] K. Hennessy, A. Badolato, M. Winger, D. Gerace, M. Atatüre, S. Gulde, S. Fält, E. L. Hu, and A. Imamoglu, *Nature* **445**, 896 (2007).
 - [6] A. Imamoglu, D. D. Awschalom, G. Burkard, D. P. DiVincenzo, D. Loss, M. Sherwin, and A. Small, *Phys. Rev. Lett.* **83**, 4204 (1999).
 - [7] R. H. Dicke, *Phys. Rev.* **93**, 99 (1954).
 - [8] M. Tavis and F. W. Cummings, *Phys. Rev.* **170**, 379 (1968).
 - [9] G. Scharf, *Helv. Phys. Acta* **43**, 806(1970).
 - [10] K. Hepp and E. H. Lieb, *Ann. Phys. (N.Y.)* **76**, 360 (1973).
 - [11] P. R. Eastham and P. B. Littlewood, *Phys. Rev. B* **64**, 235101 (2001).
 - [12] R. Bonifacio and G. Preparata, *Phys. Rev. A* **2**, 336 (1970).
 - [13] J. S. Pratt, *Phys. Rev. B* **73**, 184413 (2006).
 - [14] Independent probe of different spins is possible as they are spatially separated. For instance, size of the exciton-polariton condensates in [3] is few μm ; cloud of cold atoms in [4] has cigar shape of length that can be 100 μm .
 - [15] The Hamiltonian Eq.(6) was obtained using perturbation theory for $\epsilon \approx \omega \gg g^2/(\epsilon - \omega)$. Expressing XY term from the total spin conservation law the Hamiltonian Eq.(6) is $H = \omega b^\dagger b + \left(\epsilon + \frac{g^2(2b^\dagger b + 1)}{N(\epsilon - \omega)} \right) \sum_j S_j^z - \frac{g^2}{N(\epsilon - \omega)} \left(\sum_j S_j^z \right)^2$. Its ground state is all spin polarized up within the applicability limits of perturbation theory. The same ground state is predicted by mean-field theory in this regime.
 - [16] M. Trif, V. N. Golovach, and D. Loss, *Phys. Rev. B* **77**, 045434 (2008).
 - [17] Vladimir Buzek, Miguel Orszag, and Marian Rosko, *Phys. Rev. Lett.* **94**, 163601 (2005).
 - [18] A. Sørensen, L.-M. Duan, J. I. Cirac, and P. Zoller, *Nature* **409**, 63 (2001).
 - [19] O. Tsyplatyev and D. Loss, to appear in PRA.